

A satellite image of the Atlantic Ocean, showing a large tropical cyclone with a distinct eye and spiral cloud bands. The map includes the eastern coast of North America and the western coast of Europe. The text is overlaid on the image.

Multi-annual forecasts of Atlantic tropical cyclones in a climate service context

Louis-Philippe Caron

WWSOC, Montreal, August
2014

Objectives

- Evaluating the capability of initialized GCMs at predicting Atlantic Tropical Cyclone activity over a 5-year horizon
- Evaluating predictability of climate shifts from active to inactive regime (and vice-versa)


Two techniques are typically used to evaluate TC activity in climate simulations:

1. Direct analysis of tropical cyclones statistics
 - Low-res GCMs are bad at simulating TC activity in the Atlantic
 - Huge amount of data required
1. Indirectly through changes in large-scale fields that impact TC activity

Climate factors influencing Atlantic hurricane activity


Climate factor	Description	Timescale
Atlantic Multidecadal Oscillation (AMO or AMV)	Oscillation in North Atlantic Ocean Temp.	Decadal
El Niño Southern Oscillation	Oscillation in Tropical Pacific Ocean Temp.	Annual (~3-5 yr cycle)
West African Monsoon	Rainfall over Sahel region	Annual
North Atlantic Oscillation (NAO)	Seesaw pattern in sea level pressures b/w Iceland and the Azores	Annual
Solar activity		11-year cycle
Ozone concentration in upper atmosphere		Annual
Dust/aerosols over the Atlantic	Dust originating from Sahara desert	Annual
Madden-Julian Oscillation	Eastward propagating disturbances in the tropics	Intra-seasonal

AMO index definition:




from Klotzbach and Gray
(2008)

the 5-yr mean anomaly



Number of seasonal TC metrics above climatological mean




TC metrics are:

- Number of hurricanes
- Number of major hurricanes
- Number of hurricane days
- Accumulated Cyclone Energy (ACE)

Ensemble initialized near-term predictions


CMIP5: ensemble forecast systems using an initialized ESM




Ensemble climate forecast systems

Assume a multi-model ensemble system with coupled initialized GCMs


Model 1 Model 2 Model 3 Model 4 Model 5 Model 6




DECADAL RETROSPECTIVE FORECASTS (re-forecasts or hindcasts)




SST – Initialized Forecasts




SST – Non-Initialized Forecasts



Detrended SST – Initialized Forecasts







Detrended SST – Non-Initialized Forecasts




ACC Year 1-5



ACC, 5-year mean detrended SST – Initialized Forecasts



MSLP – Initialized Forecasts







MSLP – Non-Initialized Forecasts




ACC Year 1-5

ACC, 5-year mean MSLP – Initialized Forecasts




AMO index


Average year 1-5




ACC – 5 year mean




Correlation Obs. Index vs TC activity




Correlation Pred. Index (Ini) vs TC activity




Correlation Pred. Index (Nolni) vs TC activity




Major Hurricanes Initialized




Major Hurricanes Non-Initialized




Accumulated Cyclone Energy (ACE) - Initialized



Accumulated Cyclone Energy (ACE) – Non-Initialized







Entire period

1966-1969

1991-1994

Average AMO index
Years 2-6 minus
AMO index Year 1



Summary

- Initialized GCMs do seem capable of predicting AMO index, which is linked to Atlantic TC level, at multi-annual timescale (5yrs)
- Skill doesn't come only from persistence, i.e. we can predict shift between active and quiet phases



Thank you